### POST-SURGERY INSTRUCTIONS



It is important to take care of your surgery site, so please follow these instructions:

Bite on gauze for thirty minutes to stop any bleeding.

If bleeding continues, place another gauze and continue to apply pressure.

2 For at least 24 hours DO NOT:

Smoke	Drink through a straw	Spit or rinse excessively	Drink alcohol, carbonated liquids or very hot liquids
-------	--------------------------	---------------------------	---

Any of these can cause the blood clot over the surgery site to dislodge from the socket, causing a "dry socket" and resulting in unnecessary pain. Should this happen, call the office at 330.825.4446 to see the doctor as soon as possible.



# Maintain a soft diet:

For the remainder of the day-things like soup, ice cream, Jell-O, applesauce, etc.



#### If your pain, discomfort or bleeding continues or worsens:

Call the office 330.825.4446 for a reexamination.

Biting on a wet tea bag or an ice cube for 30 minutes can also help stop any bleeding.



## If you were given medications:

Take them according to the instructions.



### If swelling occurs:

Place an ice pack directly on your face in the area of swelling. Fifteen minutes on followed by a five-minute break repeated for several hours should reduce the swelling.



## Starting tomorrow morning

Rinse gently with warm salt water 1/4 teaspoon of salt to 1 cup of water every 4 hours for 2 days.



#### If any problems or questions do arise,

Call the office at 330.825.4446 or Dr. Kate personally at 330.760.0654.

